


1. Sketch the following vector fields:

(a) $\vec{F}_1(x, y) = y\hat{i} + \hat{j}$


(b) $\vec{F}_2(x, y) = y \cos x \hat{i} + \sin x \hat{j}$

(c) $\vec{F}_3(x, y) = -\frac{y}{\sqrt{x^2+y^2}}\hat{i} + \frac{x}{\sqrt{x^2+y^2}}\hat{j}$

Here they are, courtesy of Mathematica:



2. Give the equation defining each of the vector fields illustrated below:



The first one is a constant vector field, since all of the arrows are the same. Looking at the slope, we might guess $\vec{F}(x, y) = 2\hat{i} + \hat{j}$. (It's hard to tell for sure from the picture)

This is a vector field which at (x, y) points in the direction from the origin to (x, y) . Thus it's some multiple of $\vec{F}(x, y) = x\hat{i} + y\hat{j}$. Since all the lengths of the vectors in the picture are 1, it must be normalized so $\vec{F}(x, y) = \frac{x}{\sqrt{x^2+y^2}}\hat{i} + \frac{y}{\sqrt{x^2+y^2}}\hat{j}$.

The third is a bit tricky. The first thing to notice is that the horizontal part is constant, so the field is of the form $\vec{F}(x, y) = \hat{i} + N(x, y)\hat{j}$. Notice that $N(x, y)$ appears not to depend on x at all: for any value of x , as we change y the field changes in exactly the same way. Moreover, it seems to be periodic in y (well, this might be hard to see since I didn't include a big enough region – trust me here). The vertical part vanishes at $0, \pi/2, \pi, \dots$. So it looks like a pretty good guess is that the field is $\vec{F}(x, y) = \hat{i} + \sin(2y)\hat{j}$.

3. (4B-2) Evaluate the integral of $\vec{F} = x\hat{i} + y\hat{j}$ along the path C which goes once counter-clockwise around a circle of radius a centered at the origin. First argue geometrically, and then check your answer by directly computing the integral.

Well, $\frac{d\vec{r}}{dt}$ is always tangent to the circle (this is the velocity). But $F(\vec{r}(t))$ is a radius. The two are perpendicular, so their dot product is 0, and the line integral must be 0. It's not much harder to make this precise. We parametrize the path via $\vec{r}(t) = \langle a \cos t, a \sin t \rangle$, and so the velocity is $d\vec{r}/dt = \langle -a \sin t, a \cos t \rangle$. The vector field is $\vec{F}(\vec{r}(t)) = \langle a \cos t, a \sin t \rangle$. So

$$\int_C F d\vec{r} = \int_{t=0}^{2\pi} \langle a \cos t, a \sin t \rangle \cdot \langle -a \sin t, a \cos t \rangle dt = \int_{t=0}^{2\pi} 0 dt = 0.$$

4. For the first and second vector fields of problem 1, compute the integral along three different paths from $(0, 0)$ to $(1, 1)$:

- (a) C_1 , a straight line from $(0, 0)$ to $(1, 1)$
 (b) C_2 , a line from $(0, 0)$ to $(1, 0)$ and then to $(1, 1)$.
 (c) C_3 , along the parabola $y = x^2$

The paths are parametrized by $\vec{r}(t) = \langle t, t \rangle$, $\vec{r}(t) = \langle t, t^2 \rangle$, and $vecr(t) = \langle t, 0 \rangle$ followed by $\vec{r}(t) = \langle 1, t \rangle$. The velocity vectors are respectively $\langle 1, 1 \rangle$, $\langle 1, 2t \rangle$, and $\langle 1, 0 \rangle$ followed by $\langle 0, 1 \rangle$. For the vector field $\vec{F}(x, y) = y\hat{i} + \hat{j}$, we compute the integrals as

$$\begin{aligned} \int_{C_1} \vec{F} \cdot d\vec{r} &= \int_{t=0}^1 \langle t, 1 \rangle \cdot \langle 1, 1 \rangle dt = \int_{t=0}^1 (t + 1) dt = \frac{3}{2}, \\ \int_{C_2} \vec{F} \cdot d\vec{r} &= \int_{t=0}^1 \langle t^2, 1 \rangle \cdot \langle 1, 2t \rangle dt = \int_{t=0}^1 (t^2 + 2t) dt = \frac{4}{3}, \\ \int_{C_3} \vec{F} \cdot d\vec{r} &= \int_{t=0}^1 \langle 0, 1 \rangle \cdot \langle 1, 0 \rangle dt + \int_{t=0}^1 \langle t, 1 \rangle \cdot \langle 0, 1 \rangle dt = 0 + 1 = 1. \end{aligned}$$

For the second field, $\vec{F}(x, y) = y \cos x \hat{i} + \sin x \hat{j}$.

$$\begin{aligned} \int_{C_1} \vec{F} \cdot d\vec{r} &= \int_{t=0}^1 \langle t \cos t, \sin t \rangle \cdot \langle 1, 1 \rangle dt = \int_{t=0}^1 (t \cos t + \sin t) dt \\ &= (t \sin t)|_0^1 = \sin 1, \\ \int_{C_2} \vec{F} \cdot d\vec{r} &= \int_{t=0}^1 \langle t^2 \cos t, \sin t \rangle \cdot \langle 1, 2t \rangle dt = \int_{t=0}^1 (t^2 \cos t + 2t \sin t) dt \\ &= (t^2 \sin t)|_0^1 = \sin 1, \\ \int_{C_3} \vec{F} \cdot d\vec{r} &= \int_{t=0}^1 \langle 0, \sin t \rangle \cdot \langle 1, 0 \rangle dt + \int_{t=0}^1 \langle t \cos 1, \sin 1 \rangle \cdot \langle 0, 1 \rangle dt \\ &= 0 + \sin 1 \int_0^1 dt = \sin 1.5 \end{aligned}$$

5. One of the vector fields from the previous question is conservative. Which one? Find a function $f(x, y)$ of which it is the gradient, and evaluate the above integral using the fundamental theorem for line integrals.

We got the same integral over all three curves when integrating the second field, suggesting that it is probably a gradient field. In fact, it is the gradient field of $f(x, y) = y \sin x$, and the fundamental theorem of calculus for line integrals implies that

$$\int_C \vec{F} \cdot d\vec{r} = f(1, 1) - f(0, 0) = \sin 1 - 0 = \sin 1,$$

for any path C between $(0, 0)$ and $(1, 1)$, including the three used above.

6. What is the gradient field associated with the function $f(r, \theta) = r \log r$?

One way to do this is with the chain rule:

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{\partial f}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial f}{\partial \theta} \frac{\partial \theta}{\partial x} = (1 + \log r)(x/r) + 0 \frac{\partial \theta}{\partial x} = (1 + \log r)(x/r) \\ \frac{\partial f}{\partial y} &= \frac{\partial f}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial f}{\partial \theta} \frac{\partial \theta}{\partial y} = (1 + \log r)(y/r) + 0 \frac{\partial \theta}{\partial y} = (1 + \log r)(y/r). \end{aligned}$$

You could now plug in the usual formulas for polar coordinates to express the answer in terms of only x and y , but leaving it this way is OK too.

